


CURVAS NICE (National Collaborating Centre for Women's and Children's Health).
 The Royal College of Obstetricians and Gynaecologists. UK. 2010.

EG: 23 sem

	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	4.09	7.60
2º día	5.85	10.53
3er día y >	7.60	13.45


EG: 23 semanas



**CURVAS NICE (National Collaborating Centre for Women's and Children's Health).
The Royal College of Obstetricians and Gynaecologists. UK. 2010.**

EG: 24 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	4.09	7.89
2º día	6.43	11.11
3er día y >	8.19	14.04


EG: 24 semanas



**CURVAS NICE (National Collaborating Centre for Women's and Children's Health).
The Royal College of Obstetricians and Gynaecologists. UK. 2010.**

EG: 25 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	4.39	8.19
2º día	6.43	11.11
3er día y >	8.77	14.62


EG: 25 semanas



**CURVAS NICE (National Collaborating Centre for Women's and Children's Health
The Royal College of Obstetricians and Gynaecologists. UK. 2010.**

EG: 26 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	4.68	8.29
2º día	7.02	11.70
3er día y >	9.36	15.20


EG: 26 semanas



**CURVAS NICE (National Collaborating Centre for Women's and Children's Health).
The Royal College of Obstetricians and Gynaecologists. UK. 2010.**

EG: 27 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	4.68	8.48
2º día	7.60	11.99
3er día y >	9.94	15.79


EG: 27 semanas



RVAS NICE (National Collaborating Centre for Women's and Children's Health).
 The Royal College of Obstetricians and Gynaecologists. UK. 2010.

EG: 28 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	4.97	8.48
2º día	7.89	12.28
3er día y >	10.53	16.37


EG: 28 semanas



**CURVAS NICE (National Collaborating Centre for Women's and Children's Health).
The Royal College of Obstetricians and Gynaecologists. UK. 2010.**

EG: 29 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	5.26	8.77
2º día	8.19	12.87
3er día y >	11.11	16.96


EG: 29 semanas



**CURVAS NICE (National Collaborating Centre for Women's and Children's Health).
The Royal College of Obstetricians and Gynaecologists. UK. 2010.**

EG: 31 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	5.56	9.06
2º día	8.77	13.74
3er día y >	12.28	18.13


EG: 31 semanas



**CURVAS NICE (National Collaborating Centre for Women's and Children's Health).
The Royal College of Obstetricians and Gynaecologists. UK. 2010.**

EG: 32 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	5.85	9.36
2º día	9.36	14.04
3er día y >	12.87	18.71


EG: 32 semanas



**CURVAS NICE (National Collaborating Centre for Women's and Children's Health).
The Royal College of Obstetricians and Gynaecologists. UK. 2010.**

EG: 33 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	6.14	9.65
2º día	9.94	14.33
3er día y >	13.45	19.30


EG: 33 semanas



**CURVAS NICE(National Collaborating Centre for Women's and Children's Health).
The Royal College of Obstetricians and Gynaecologists. UK. 2010.**

EG: 34 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	6.43	9.94
2º día	10.23	14.62
3er día y >	14.04	19.88

EG: 34 semanas



CURVAS NICE (National Collaborating Centre for Women's and Children's Health).
 The Royal College of Obstetricians and Gynaecologists. UK. 2010.

EG: 35 sem	Fototerapia (mg/dl)	Exanguíneo Transfusión (mg/dl)
1er día	6.73	10.23
2º día	10.53	15.20
3er día y >	14.62	20.47

EG: 35 semanas

